



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

3. Proposed by CHARLES E. MYERS, Canton, Ohio.

A spherical air-bubble, having risen from a depth of 1,500 feet in water, was one inch in diameter when it reached the surface; what was its diameter at the point of starting?

4. Proposed by DeVOLSON WOOD, M.A., C. E., Professor of Mechanical Engineering, Stevens Institute of Technology, Hoboken, New Jersey.

A particle starts at rest and revolves in a circle with a uniform acceleration, acquiring a velocity v in t seconds. Find the locus of the foot of the perpendicular from the centre of the circle upon the resultant acceleration.

5. Proposed by J. B. BALDWIN, A. M., Professor of Mathematics and Commercial Law, Davenport Business College, Davenport, Iowa.

A 200 pound ball lies on a three legged table, having the legs equally distant apart and perpendicular to the plane of the top of the table. (1) What is the weight on each leg of the table not including the top when the ball is 2 feet, 3 feet, and 4 feet distant from the three legs? (2) If the ball is 2 feet, 3 feet, and 5 feet from the legs, what must be the weight of the top to keep from tipping and the weight on each leg excluding the top and also including the top?

DIOPHANTINE ANALYSIS.

Conducted by J. M. COLAW, Monterey, Va. All contributions to this department should be sent to him.

PROBLEMS.

1. Proposed by EARL D. WEST, West Middleburg, Logan County, Ohio.

It is required to divide a given square number into two such parts that each part will be a square number.

2. Proposed by J. M. COLAW, Principal of High School, Monterey, Virginia.

Find two numbers, such that the difference of their squares may be a cube, and the difference of their cubes a square.

3. Proposed by O. S. KIBLER, Superintendent of Schools, West Middleburg, Logan County, Ohio.

It is required to find three whole numbers in an arithmetical progression, such that the sum of every two of them shall be a square.

4. Proposed by H. W. HOLYCROSS, Superintendent of Schools, Pottersburg, Union County, Ohio.

What value of x will render $4x^4 + 12x^3 - 3x^2 - 2x + 1$ a square?

AVERAGE AND PROBABILITY.

Conducted by B. F. FINKEL, Kidder, Missouri. All contributions to this department should be sent to him.

PROBLEMS.